



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

Since for a given circle and a given exterior point the product of any secant through the point and the exterior segment of the secant is equal to the square of the tangent from the point to the circle we have  $A'B \cdot AB = x_2^2 + y_2^2 - k^2$  and  $A'B \cdot A'B' = (x_3 - k')^2 + y_3^2 - k^2$ . Hence,

$$AB + A'B' = \frac{x_2^2 + y_2^2 - k^2 + (x_3 - k')^2 + y_3^2 - k^2}{A'B}.$$

By use of  $A'B = \sqrt{(x_2 - x_3)^2 + (y_2 - y_3)^2}$  and the relations given above this equation readily reduces to  $AB + A'B' = 2k'/\sqrt{1 + m^2}$ , which is independent of  $k$  and dependent upon  $k'$  and  $\theta$  only, as was to be shown.

Also solved by T. M. BLAKSLEE, PAUL CAPRON, OLIVE C. HAZLETT, C. N. MILLS, E. J. OGLESBY, ARTHUR PELLETIER, C. P. SOUSLEY, C. E. STROMQUIST, and H. S. UHLER.

**2804 [1920, 32]. Proposed by T. H. GRONWALL, Washington, D. C.**

Show that for  $|x| < 1$

$$\begin{aligned} \frac{1}{\sqrt{1-x^4}} \int_0^x \frac{dx}{\sqrt{1-x^4}} &= x + \sum_1^{\infty} \frac{3 \cdot 7 \cdots (4n-5)(4n-1)}{5 \cdot 9 \cdots (4n-3)(4n+1)} x^{4n+1}, \\ \left( \int_0^x \frac{dx}{\sqrt{1-x^4}} \right)^2 &= x^2 + \sum_1^{\infty} \frac{3 \cdot 7 \cdots (4n-5)(4n-1)}{5 \cdot 9 \cdots (4n-3)(4n+1)} \frac{x^{4n+2}}{2n+1}. \end{aligned}$$

SOLUTION BY O. S. ADAMS, Coast and Geodetic Survey.

The general forms for such integrals may be derived and the formulas in the statement will then be special cases of the derived forms. We then state the problem as follows:

For  $|x| < 1$ ,  $s$  a positive integer,

$$\begin{aligned} \frac{1}{\sqrt{1-x^s}} \int_0^x \frac{dx}{\sqrt{1-x^s}} &= x + \sum_1^{\infty} \frac{\frac{s+2}{2} \cdot \frac{3s+2}{2} \cdots \frac{(2n-1)s+2}{2}}{(s+1)(2s+1) \cdots (ns+1)} x^{ns+1}, \\ \left( \int_0^x \frac{dx}{\sqrt{1-x^s}} \right)^2 &= x^2 + \sum_1^{\infty} \frac{\frac{s+2}{2} \cdot \frac{3s+2}{2} \cdots \frac{(2n-1)s+2}{2}}{(s+1)(2s+1) \cdots (ns+1)} \frac{2x^{ns+2}}{ns+2}. \end{aligned}$$

Let

$$\phi(x) = \int_0^x \frac{dx}{\sqrt{1-x^s}};$$

then

$$\frac{d}{dx} \phi(x) = \frac{1}{\sqrt{1-x^s}},$$

and

$$2\phi(x) \frac{d}{dx} \phi(x) = \frac{2}{\sqrt{1-x^s}} \int_0^x \frac{dx}{\sqrt{1-x^s}}.$$

By integration,

$$\phi^2(x) = 2 \int_0^x \left[ \frac{1}{\sqrt{1-x^s}} \int_0^x \frac{dx}{\sqrt{1-x^s}} \right] dx.$$

Therefore, if the first series is multiplied by  $2dx$  and integrated from 0 to  $x$ , the second series will be obtained.

Let

$$y = \frac{1}{\sqrt{1-x^s}} \int_0^x \frac{dx}{\sqrt{1-x^s}};$$

then

$$\frac{dy}{dx} = \frac{1}{1-x^s} + \frac{(s/2)x^{s-1}}{(1-x^s)^{3/2}} \int_0^x \frac{dx}{\sqrt{1-x^s}} = \frac{1}{1-x^s} + \frac{(s/2)x^{s-1}y}{1-x^s};$$

or

$$(1 - x^s) \frac{dy}{dx} - \frac{s}{2} x^{s-1} y - 1 = 0.$$

The function  $1/\sqrt{1 - x^s}$  can be developed as a power series with unit radius of convergence, this can be integrated term by term, and the result multiplied by the original series, giving for  $y$  an expression of the form  $y = x + ax^{s+1} + bx^{2s+1} + cx^{3s+1} + dx^{4s+1} + \dots$ . Now the series is absolutely convergent and we can determine the coefficients  $a, b, c$ , by substituting this series in the differential equation. We obtain

$$\begin{aligned} 1 + (s+1)ax^s + (2s+1)bx^{2s} + (3s+1)cx^{3s} + (4s+1)dx^{4s} + \dots \\ - x^s - (s+1)ax^{2s} - (2s+1)bx^{3s} - (3s+1)cx^{4s} - \dots \\ - 1 - \frac{s}{2}x^s - \frac{s}{2}ax^{2s} - \frac{s}{2}bx^{3s} - \frac{s}{2}cx^{4s} - \dots \equiv 0; \end{aligned}$$

so that the coefficients of the various powers of  $x$  must be identically zero. We must then have

$$a = \frac{\frac{s+2}{2}}{s+1}, \quad b = \frac{\frac{s+2}{2} \cdot \frac{3s+2}{2}}{(s+1)(2s+1)},$$

and

$$c = \frac{\frac{s+2}{2} \cdot \frac{3s+2}{2} \cdot \frac{5s+2}{2}}{(s+1)(2s+1)(3s+1)}.$$

The  $(n+1)$ th coefficient thus becomes equal to the expression

$$\frac{\frac{s+2}{2} \cdot \frac{3s+2}{2} \dots \frac{(2n-1)s+2}{2}}{(s+1)(2s+1) \dots (ns+1)}.$$

This proves the formula

$$\frac{1}{\sqrt{1-x^s}} \int_0^x \frac{dx}{\sqrt{1-x^s}} = x + \sum_1^\infty \frac{\frac{s+2}{2} \cdot \frac{3s+2}{2} \dots \frac{(2n-1)s+2}{2}}{(s+1)(2s+1) \dots (ns+1)} x^{ns+1}.$$

By multiplying this series by  $2dx$  and integrating from 0 to  $x$ , we obtain the second formula

$$\left( \int_0^x \frac{dx}{\sqrt{1-x^s}} \right)^2 = x^2 + \sum_1^\infty \frac{\frac{s+2}{2} \cdot \frac{3s+2}{2} \dots \frac{(2n-1)s+2}{2}}{(s+1)(2s+1) \dots (ns+1)} \frac{2x^{ns+2}}{ns+2}.$$

These formulas can easily be verified for  $s=1$  and they can be compared with the known forms for  $s=2$ . When  $s=4$ , they become identical with those stated in the problem.

$$\begin{aligned} \frac{1}{\sqrt{1-x^4}} \int_0^x \frac{dx}{\sqrt{1-x^4}} &= x + \sum_1^\infty \frac{3 \cdot 7 \dots (4n-5)(4n-1)}{5 \cdot 9 \dots (4n-3)(4n+1)} x^{4n+1}, \\ \left( \int_0^x \frac{dx}{\sqrt{1-x^4}} \right)^2 &= x^2 + \sum_1^\infty \frac{3 \cdot 7 \dots (4n-5)(4n-1)}{5 \cdot 9 \dots (4n-3)(4n+1)} \frac{x^{4n+2}}{2n+1}. \end{aligned}$$

The last formula is given by Gauss, *Werke*, vol. 3, p. 406.

Also solved by ELIJAH SWIFT and the Proposer.